



Contents lists available at ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra



The isolated quantum group of the GL_2 family



Serge Skryabin¹

*Institute of Mathematics and Mechanics, Kazan Federal University, Kremlevskaya
St. 18, 420008 Kazan, Russia*

ARTICLE INFO

Article history:

Received 6 April 2017

Available online 24 July 2017

Communicated by Michel Van den Bergh

MSC:

16T20

Keywords:

Quantum groups

Hopf algebras

ABSTRACT

Quantum analogs of $GL(2)$ are parameterized by $PGL(2)$ -orbits in the commuting variety of the projective linear group $PGL(2)$. The quantum group corresponding to the orbit constituting the second connected component of the commuting variety has previously received little attention. This paper investigates ring-theoretical properties of its coordinate ring.

© 2017 Elsevier Inc. All rights reserved.

Introduction

A basic idea going back to Manin [18] is to interpret a quantum group of the general linear type in terms of the Hopf algebra H universally coacting on a pair of quadratic graded algebras. It can be assumed that the two graded algebras have the same vector space V as the degree 1 homogeneous component generating each of them, and then the defining relations in the algebras are given by 2 vector subspaces $U, U' \subset V^{\otimes 2}$. These algebras are regarded as the quantum symmetric algebra and the quantum Grassmann algebra, and one expects that their Hilbert series should be exactly the same as in the

E-mail address: Serge.Skryabin@kpfu.ru.

¹ This work was funded by the subsidy allocated to Kazan Federal University for the state assignment in the sphere of scientific activities.